PRE-PROCEDURE INFORMATION FOR MICROPIGMENTATION

- All micropigmentation (Permanent makeup) procedures are multi-session processes. You are
 required to come back for at least one touch-up visit before it can be determined that your work is
 complete. Touch up visits are scheduled at 4 week to 3 month intervals.
- Be prepared for the color intensity of your procedure to be significantly larger, sharper, brighter or
 darker than what is expected for the final outcome. It will take time for the color to become softer
 and lighter. This transition in intensity is based on how quickly the outer layer of your skin
 softens.
- Initially, the pigment color may simulate the exact color and tone desired. Remember, it will not always remain a perfect match. Pigment colors placed in skin are constant, but your own skin tones will vary depending on exposure to cold, heat, sun, and circulatory changes. For example, if you suntan a camouflaged scar, the surrounding skin will appear darker than the treated area.
- Since delicate skin or sensitive areas may swell slightly or redden, some clients feel it best not to make any social plans for a day or two following any procedure.
- On the day of your procedure, come into the spa with no makeup.
- Any tweezing or waxing shoud be done at least 48 hours prior to procedure; electrolysis no less
 than 5 days before. Do not resume any method of hair removal for at least 2 weeks after
 procedure.
- Any eyelash or eyebrow tinting or eyelash curling should be done no sooner that 48 hours before and 2 weeks after the procedure.
- Do not wear contact lenses during or immediately following the eyeliner procedure. Remember to bring your eyeglasses. You may resume wearing your contact lenses as soon as your eyes return to their original condition before the procedure.
- If you are having lip procedures, moisturize your lips for at least 1 week before the procedure. If you have any history of cold sores/ fever blisters/ herpes simplex you will be required to contact your physician to obtain the proper prescriptive medication to help prevent such outbreaks.
- If you are taking sedatives or any medication that causes drowsiness, have someone available to drive you home.
- Refrain from the use of alcohol, aspirin, aspirin containing medications, ibuprofen or coumadin
 for 7 days before and 2 days after any procedure. No medication should be discontinued
 without first consulting your physician.

Client signature:	 Date:
_	